



ISSN 2105-2301 (supplément hebdomadaire)

**Saint-Pierre, Friday 26 February 2021, 12h (UTC -4)**

## **Weekly report on the activity of la Montagne Pelée for the period from 19<sup>th</sup> to 26<sup>th</sup> February 2021**

The seismicity of volcanic origin has increased in the last week. None of these earthquakes were felt by the population. Between February 19, 2021 at 4 p.m. UT (local time + 4 hours) and February 26, 2021 at 4 p.m. UTC (local time + 4 hours), the OVSM recorded at least 46 volcano-tectonic earthquakes of magnitude less or equal to 1, with one earthquake located at a depth of 15.1 km below sea level and the others located between 1.2 km below sea level and 1 km above sea level. The shallow depth volcano-tectonic type seismicity, characterized by high frequency signals, is associated with micro-fracturing in the volcanic edifice whereas the deep volcano-tectonic type earthquake may be related to the pressurization of a deep zone of the magmatic system.

During the period of this report, OVSM recorded 1 isolated long-period type earthquake. This type of signal containing low frequencies is associated with the migration of fluids in the volcanic edifice.

During phases of volcanic reactivation, it is common for periods of increased seismic activity to alternate with phases of lower seismic activity. The seismicity of volcanic origin remains above the average base level recorded between January 1, 2015 and April 2019 (the date of the start of its increase).

A zone of strongly degraded, browned and dead vegetation is still observed on the southwest flank of Mount Pelée, between the upper Claire River and the Chaude River, above their confluence. A field expedition on the ground on February 8, 2021 by the OVSM IPGP, with the support of the STIS and the civil protection helicopter, confirmed the presence in this zone of diffuse and passive soil degassing, without the presence of fumaroles, of carbon dioxide (CO<sub>2</sub>, an odorless and colorless gas) significantly higher than the base level and which was measured with specific and calibrated detectors of OVSM. This area is located less than 200 m from the Claire River where the presence of hydrogen sulfide (H<sub>2</sub>S, smell of rotten egg) has been reported in scientific literature for several years, in the vicinity of thermal springs that originate from the hydrothermal system of the Mount Pelée.

This report and the monthly OVSM-IPGP bulletins are available at <http://www.ipgp.fr/fr/ovsm/actualites-communiqués-publics-de-lovsm>. You can also find us on our Facebook and Twitter accounts.

The alert level remains yellow: vigilance.

The Director's office of OVSM IPGP.